

MAT 167: STATISTICS

TEST I: CHAPTERS 1-4

INSTRUCTOR: ANTHONY TANBAKUCHI

SPRING 2008

Name: _____

Computer / Seat Number: _____

No books, notes, or friends. **Show your work.** You may use the attached equation sheet, R, and a calculator. No other materials. Write your work in the provided space for each problem (including any R work if appropriate). You may not use personal computers, only use the classroom computer at your desk. Using any other program or having any other documents open on the computer will constitute cheating.

You have until the end of class to finish the exam, manage your time wisely.

If something is unclear quietly come up and ask me.

If the question is legitimate I will inform the whole class.

Express all final answers to 3 significant digits. Probabilities should be given as a decimal number unless a percent is requested. Circle final answers, ambiguous or multiple answers will not be accepted. Show steps where appropriate.

The exam consists of 10 questions for a total of 46 points on 7 pages.

This Exam is being given under the guidelines of our institution's **Code of Academic Ethics**. You are expected to respect those guidelines.

Points Earned: _____ out of 46 total points

Exam Score: _____

1. Provide **short succinct** written answers to the following conceptual questions.

- (a) (1 point) Would mass in grams be classified as a nominal, ordinal, interval, or ratio level of measurement?
- (b) (1 point) Which of the following measures of variation is most effected by outliers:
IQR, standard deviation, range
- (c) (1 point) What percent of data is greater than Q_3 ?
- (d) (1 point) If the mean, median, and mode for a data set are all the same, what can you conclude about the data's distribution?
- (e) (1 point) If the median is greater than the mode for a data set, what can you conclude about the data's distribution?
- (f) (1 point) What does the standard deviation represent conceptually **in words**? (Be concise but don't simply state the equation in words verbatim.)
- (g) (1 point) Give an example of sampling error?
- (h) (2 points) A box plot is a useful tool that can quickly communicate many traits about a set of data. List 4 useful pieces of information that an observer can easily assess using a box plot.
- (i) (1 point) You scored in the 78th percentile on the GRE. If 8,000 people took the GRE, how many people had a score at least as high as your score?

2. A survey conducted in our class asked 18 students how many hours they work per week. Use the R output below to answer the following questions.

There are 18 data points stored in the variable x , below is the sorted data:

```
> sort(x)
[1] 0 0 0 0 8 8 10 12 13 20 21 30 30 30 30 35 50
```


The basic descriptive statistical analysis is as follows:

```
> summary(x)
  Min. 1st Qu.  Median    Mean 3rd Qu.    Max.
  0.00   8.00   16.50   18.17   30.00   50.00

> var(x)
[1] 215.6765

> sd(x)
[1] 14.68593

> hist(x, xlab = "hours", main = "Student Enrollment Data")
```




- (a) (1 point) Use the range rule of thumb to estimate the standard deviation. Is it close to the actual standard deviation?
- (b) (1 point) What is P_{50} equal to?
- (c) (1 point) What is the IQR (inter quartile range) equal to?

- (d) (1 point) What percent of the data falls within the IQR?
- (e) (1 point) What is the mode for the data?
- (f) (1 point) What is the percentile for a student who works 8 hours per week?
- (g) (1 point) What is the z -score for the student who is working 50 hours per week?
- (h) (1 point) Is 35 hours an unusual (outlier) value based on its z score? (Why)
- (i) (2 points) If someone asked what the typical number of hours worked per week for this class was, why would just reporting the mean be misleading? **What would you report in addition to the mean?**
- (j) (1 point) Is the data positively skewed, negatively skewed, or symmetrical?
- (k) (1 point) Construct an interval using the Empirical Rule which you would expect 95% of the data to fall within.

(1) (1 point) Would the Empirical Rule be accurate when used for this data set? **Why?**

3. (1 point) When plotting frequencies tabulated from categorical data, why would a bar graph be preferable to a pie chart?

4. Use the below box plot to answer the following questions.



(a) (1 point) Which gender has a higher median number of credits?

(b) (1 point) What is the approximate median number of credits for the females?

(c) (1 point) What is the approximate IQR of credits for the males?

(d) (1 point) Which gender has a larger range of credits?

(e) (1 point) How many outliers are there in this data set as indicated by the box plots?

5. Using the below table for our class to answer the following questions.

	BLACK	BLOND	BROWN
CAR	0	2	11
PUBLIC	2	0	0
TRUCK	1	0	2

- (a) (1 point) Find the probability of selecting a person with brown hair.
- (b) (1 point) Would it be unusual to randomly select a person with brown hair?
- (c) (1 point) Find the probability of randomly selecting three car drivers without replacement.
- (d) (1 point) If you randomly select 4 people with replacement, what is the probability that at least one uses public transportation?
- (e) (1 point) Find the probability of selecting a student who drives a truck or a student with brown hair.
- (f) (1 point) Find the probability of selecting a person with blond hair given that they are a truck driver.

6. A researcher needs to estimate the mean height of corn in a field containing 10,000 corn plants. The corn plants are planted 100 to a row, and there are 100 rows. The researcher wants to take a sample of 100 corn plant heights.
- (a) (1 point) Describe a method for sampling this population that would be classified as a convenience sample.

 - (b) (1 point) Describe a method for sampling this population that would be classified as a simple random sample.
7. The weather report for this work week (Monday through Friday) states that the probability of rain is 5% for each day.
- (a) (1 point) What is the probability that it will rain at least once this week?

 - (b) (1 point) What is the probability that it won't rain this week?

 - (c) (1 point) If you decided not to use an umbrella this week, would it be unusual for you to get wet?

8. (2 points) Car tires must not deform or explode when inflated up to their maximum pressure rating. Before distributing the tires, they must be tested. To test the safety of tires, an inspector randomly samples 50 tires (without replacement) from a batch of 5,000 that have been manufactured. The inspector inflates each of the fifty tires until they explode or deform to make sure they meet the minimum safety requirements. If none of the sampled tires fails the test, the tires will be distributed to dealers. If the batch contains 15 defective tires that will explode if selected, what is the probability that the batch will be rejected?

9. (2 points) The quadratic mean is defined as

$$\text{quadratic mean} = \sqrt{\frac{\sum x_i^2}{n}}$$

find the quadratic mean of x . If $x = \{2, 8, 7, 4, 9\}$

10. (2 points) Given $y = \{a, -2a, 4a\}$, completely simplify the following expression. Assume a is an unknown constant.

$$\frac{(\sum (y_i - 2a))^2}{9a}$$

Statistics Quick Reference

Card & R Commands

by Anthony Tanbakuchi. Version 1.8.2
<http://www.tanbakuchi.com>
 ANTHONY@TANBAKUCHI.COM
 Get R at: <http://www.r-project.org>
 R commands: bold typewriter text

1 Misc R

To make a vector v store data: $x=c(x1, x2, \dots)$
 Help: general `RSiteSearch("Search Phrase")`
 Get: function `?functionName`
 Get column of data from table:
`tableName$columnName`
 List all variables: `ls()`
 Delete all variables: `rm(list=ls())`

$$\sqrt{x} = \text{sqrt}(x) \quad (1)$$

$$x^n = x^n \quad (2)$$

$$n = \text{length}(x) \quad (3)$$

$$T = \text{table}(x) \quad (4)$$

2 Descriptive Statistics

2.1 NUMERICAL

Let $x=c(x1, x2, x3, \dots)$

$$\text{total} = \sum_{i=1}^n x_i = \text{sum}(x) \quad (5)$$

$$\text{min} = \text{min}(x) \quad (6)$$

$$\text{max} = \text{max}(x) \quad (7)$$

$$\text{six number summary} = \text{summary}(x) \quad (8)$$

$$\mu = \frac{\sum x_i}{N} = \text{mean}(x) \quad (9)$$

$$\bar{x} = \frac{\sum x_i}{N} = \text{mean}(x) \quad (10)$$

$$\bar{x} = P_{50} = \text{median}(x) \quad (11)$$

$$\sigma = \sqrt{\frac{\sum (x_i - \mu)^2}{N}} \quad (12)$$

$$s = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n-1}} = \text{sd}(x) \quad (13)$$

$$CV = \frac{\sigma}{\mu} = \frac{s}{\bar{x}} \quad (14)$$

2.2 RELATIVE STANDING

$$z = \frac{x - \mu}{\sigma} = \frac{x - \bar{x}}{s} \quad (15)$$

Percentiles:

$$P_k = x_i, \text{ (sorted } x) \quad (16)$$

$$k = \frac{i-0.5}{n} \cdot 100\%$$

To find x_i given P_k , i is:

- $L = (k/100)n$
- if L is an integer: $i = L + 0.5$; otherwise $i = L$ and round up.

2.3 VISUAL

All plots have optional arguments:

- `main=""` sets title
- `xlab=""`, `ylab=""` sets x/y-axis label
- `type="p"` for point plot
- `type="l"` for line plot
- `type="b"` for both points and lines

Ex: plot(x, y, type="b", main="My Plot")

Plot Types:

- `hist(x)` histogram
- `stem(x)` stem & leaf
- `boxplot(x)` box plot
- `plot(T)` bar plot, `T=table(x)`
- `plot(x, y)` scatter plot, x, y are ordered vectors
- `plot(t, y)` time series plot, t, y are ordered vectors
- `curve(expr, xmin, xmax)` plot expr involving x

2.4 ASSESSING NORMALITY

Q-Q plot: `qqnorm(x)`; `qqline(x)`

3 Probability

Number of successes x with n possible outcomes. (Don't double count!)

$$P(A) = \frac{x}{n} \quad (17)$$

$$P(\bar{A}) = 1 - P(A) \quad (18)$$

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B) \quad (19)$$

$$P(A \text{ or } B) = P(A) + P(B) \text{ if } A, B \text{ mut. excl.} \quad (20)$$

$$P(A \text{ and } B) = P(A) \cdot P(B|A) \quad (21)$$

$$P(A \text{ and } B) = P(A) \cdot P(B) \text{ if } A, B \text{ independent} \quad (22)$$

$$n! = n(n-1) \cdots 1 = \text{factorial}(n) \quad (23)$$

$${}_n P_k = \frac{n!}{(n-k)!} \text{ Perm. no elem. alike} \quad (24)$$

$${}_n C_k = \frac{n!}{n_1! n_2! \cdots n_k!} \text{ Perm. } n_1 \text{ alike, } \dots \quad (25)$$

$${}_n C_k = \frac{n!}{(n-k)! k!} = \text{choose}(n, k) \quad (26)$$

4 Discrete Random Variables

$$P(x_i) : \text{probability distribution} \quad (27)$$

$$E = \mu = \sum x_i \cdot P(x_i) \quad (28)$$

$$\sigma = \sqrt{\sum (x_i - \mu)^2 \cdot P(x_i)} \quad (29)$$

4.1 BINOMIAL DISTRIBUTION

$$\mu = n \cdot p \quad (30)$$

$$\sigma = \sqrt{n \cdot p \cdot q} \quad (31)$$

$$P(x) = {}_n C_x p^x q^{n-x} = \text{dbinom}(x, n, p) \quad (32)$$

4.2 POISSON DISTRIBUTION

$$P(x) = \frac{\mu^x \cdot e^{-\mu}}{x!} = \text{dpois}(x, \mu) \quad (33)$$

5 Continuous random variables

CDF $F(x)$ gives area to the left of x , $F^{-1}(p)$ expects p is area to the left.

$$f(x) : \text{probability density} \quad (34)$$

$$E = \mu = \int_{-\infty}^{\infty} x \cdot f(x) dx \quad (35)$$

$$\sigma = \sqrt{\int_{-\infty}^{\infty} (x - \mu)^2 \cdot f(x) dx} \quad (36)$$

$$F(x) : \text{cumulative prob. density (CDF)} \quad (37)$$

$$F^{-1}(x) : \text{inv. cumulative prob. density} \quad (38)$$

$$F(x) = \int_{-\infty}^x f(x') dx' \quad (39)$$

$$p = P(x < x') = F(x') \quad (40)$$

$$x' = F^{-1}(p) \quad (41)$$

$$p = P(x > a) = 1 - F(a) \quad (42)$$

$$p = P(a < x < b) = F(b) - F(a) \quad (43)$$

5.1 UNIFORM DISTRIBUTION

$$p = P(a < u') = F(u') \quad (44)$$

$$u' = F^{-1}(p) = \text{qunif}(p, \text{min}=a, \text{max}=b) \quad (45)$$

5.2 NORMAL DISTRIBUTION

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2\sigma^2} \frac{(x-\mu)^2}{\sigma^2}} \quad (46)$$

$$p = P(z < z') = F(z') = \text{pnorm}(z') \quad (47)$$

$$z' = F^{-1}(p) = \text{qnorm}(p) \quad (48)$$

$$p = P(x < x') = F(x') \quad (49)$$

$$= \text{pnorm}(x', \text{mean}=\mu, \text{sd}=\sigma) \quad (49)$$

$$x' = F^{-1}(p) \quad (50)$$

$$= \text{qnorm}(p, \text{mean}=\mu, \text{sd}=\sigma) \quad (50)$$

5.3 t-DISTRIBUTION

$$p = P(t < t') = F(t') = \text{pt}(t', \text{df}) \quad (51)$$

$$t' = F^{-1}(p) = \text{qt}(p, \text{df}) \quad (52)$$

5.4 χ^2 -DISTRIBUTION

$$p = P(\chi^2 < \chi'^2) = F(\chi'^2) \quad (53)$$

$$= \text{pchisq}(\chi'^2, \text{df}) \quad (53)$$

$$\chi'^2 = F^{-1}(p) = \text{qchisq}(p, \text{df}) \quad (54)$$

5.5 F-DISTRIBUTION

$$p = P(F < F') = F(F') \quad (55)$$

$$= \text{pf}(F', \text{df1}, \text{df2}) \quad (55)$$

$$F' = F^{-1}(p) = \text{qf}(p, \text{df1}, \text{df2}) \quad (56)$$

6 Sampling distributions

$$\mu_{\bar{x}} = \mu \quad \sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}} \quad (57)$$

$$\mu_{\bar{p}} = p \quad \sigma_{\bar{p}} = \sqrt{\frac{pq}{n}} \quad (58)$$

7 Estimation

7.1 CONFIDENCE INTERVALS

$$\text{proportion: } \hat{p} \pm E, \quad E = z_{\alpha/2} \cdot \sigma_{\bar{p}} \quad (59)$$

$$\text{mean } (\sigma \text{ known): } \bar{x} \pm E, \quad E = z_{\alpha/2} \cdot \sigma_{\bar{x}} \quad (60)$$

$$\text{mean } (\sigma \text{ unknown, use } s): \bar{x} \pm E, \quad E = t_{\alpha/2} \cdot \sigma_{\bar{x}} \quad (61)$$

$$df = n - 1$$

$$\text{variance: } \frac{(n-1)s^2}{\chi^2_R} < \sigma^2 < \frac{(n-1)s^2}{\chi^2_L} \quad (62)$$

$$df = n - 1$$

$$2 \text{ proportions: } \Delta \hat{p} \pm z_{\alpha/2} \cdot \sqrt{\frac{\hat{p}_1 \hat{q}_1}{n_1} + \frac{\hat{p}_2 \hat{q}_2}{n_2}} \quad (63)$$

$$2 \text{ means (indep): } \Delta \bar{x} \pm t_{\alpha/2} \cdot \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}} \quad (64)$$

$$df \approx \text{min}(n_1 - 1, n_2 - 1)$$

$$\text{matched pairs: } \bar{d} \pm t_{\alpha/2} \cdot \frac{s_d}{\sqrt{n}}, \quad d_i = x_i - y_i, \quad (65)$$

$$df = n - 1$$

7.2 CI CRITICAL VALUES (TWO SIDED)

$$z_{\alpha/2} = F_z^{-1}(1 - \alpha/2) = \text{qnorm}(1 - \alpha/2) \quad (66)$$

$$t_{\alpha/2} = F_t^{-1}(1 - \alpha/2) = \text{qt}(1 - \alpha/2, \text{df}) \quad (67)$$

$$\chi^2_L = F_{\chi^2}^{-1}(\alpha/2) = \text{qchisq}(\alpha/2, \text{df}) \quad (68)$$

$$\chi^2_R = F_{\chi^2}^{-1}(1 - \alpha/2) = \text{qchisq}(1 - \alpha/2, \text{df}) \quad (69)$$

7.3 REQUIRED SAMPLE SIZE

$$\text{proportion: } n = \hat{p} \hat{q} \left(\frac{z_{\alpha/2}}{E} \right)^2, \quad (70)$$

$$(\hat{p} = \hat{q} = 0.5 \text{ if unknown})$$

$$\text{mean: } n = \left(\frac{z_{\alpha/2} \cdot \hat{\sigma}}{E} \right)^2 \quad (71)$$

8 Hypothesis Tests

Test statistic and R function (when available) are listed for each.

Optional arguments for hypothesis tests:

alternative="two.sided" can be:
"two.sided", "less", "greater"

conf.level=0.95 constructs a 95% confidence interval. Standard CI only when **alternative**="two.sided".

Optional arguments for power calculations & Type II error:

alternative="two.sided" can be:
"two.sided" or "one.sided"

sig.level=0.05 sets the significance level α .

8.1 1-SAMPLE PROPORTION

$H_0: p = p_0$

prop.test(x, n, p=p₀, alternative="two.sided")

$$z = \frac{\hat{p} - p_0}{\sqrt{p_0 q_0 / n}} \quad (72)$$

8.2 1-SAMPLE MEAN (σ KNOWN)

$H_0: \mu = \mu_0$

$$z = \frac{\bar{x} - \mu_0}{\sigma / \sqrt{n}} \quad (73)$$

8.3 1-SAMPLE MEAN (σ UNKNOWN)

$H_0: \mu = \mu_0$

t.test(x, mu= μ_0 , alternative="two.sided")

Where **x** is a vector of sample data.

$$t = \frac{\bar{x} - \mu_0}{s / \sqrt{n}}, \quad df = n - 1 \quad (74)$$

Required Sample size:

power.t.test(delta= h , sd= σ , sig.level= α , power=1 - β , type="one.sample", alternative="two.sided")

8.4 2-SAMPLE PROPORTION TEST

$H_0: p_1 = p_2$ or equivalently $H_0: \Delta p = 0$

prop.test(x, n, alternative="two.sided")

where: **x**=c(x_1, x_2) and **n**=c(n_1, n_2)

$$z = \frac{\hat{\Delta p} - \Delta p_0}{\sqrt{\frac{\hat{p}_1 \hat{q}_1}{n_1} + \frac{\hat{p}_2 \hat{q}_2}{n_2}}}, \quad \hat{\Delta p} = \hat{p}_1 - \hat{p}_2 \quad (75)$$

$$\hat{p} = \frac{x_1 + x_2}{n_1 + n_2}, \quad \hat{q} = 1 - \hat{p} \quad (76)$$

Required Sample size:

power.prop.test(p1= p_1 , p2= p_2 , power=1 - β , sig.level= α , alternative="two.sided")

8.5 2-SAMPLE MEAN TEST

$H_0: \mu_1 = \mu_2$ or equivalently $H_0: \Delta \mu = 0$

t.test(x1, x2, alternative="two.sided")

where: **x1** and **x2** are vectors of sample 1 and sample 2 data.

$$t = \frac{\Delta \bar{x} - \Delta \mu_0}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}, \quad df \approx \min(n_1 - 1, n_2 - 1), \quad \Delta \bar{x} = \bar{x}_1 - \bar{x}_2 \quad (77)$$

Required Sample size:

power.t.test(delta= h , sd= σ , sig.level= α , power=1 - β , type="two.sample", alternative="two.sided")

8.6 2-SAMPLE MATCHED PAIRS TEST

$H_0: \mu_d = 0$

t.test(x, y, paired=TRUE, alternative="two.sided")

where: **x** and **y** are ordered vectors of sample 1 and sample 2 data.

$$t = \frac{\bar{d} - \mu_{d0}}{s_d / \sqrt{n}}, \quad d_i = x_i - y_i, \quad df = n - 1 \quad (78)$$

Required Sample size:

power.t.test(delta= h , sd= σ , sig.level= α , power=1 - β , type="paired", alternative="two.sided")

8.7 TEST OF HOMOGENEITY, TEST OF INDEPENDENCE

$H_0: p_1 = p_2 = \dots = p_k$ (homogeneity)

$H_0: X$ and Y are independent (independence)

chisq.test(D)

Enter table: **D**=data.frame(**c1, c2, ...**), where **c1, c2, ...** are column data vectors.

Or generate table: **D**=table(**x1, x2**), where **x1, x2** are ordered vectors of raw categorical data.

$$\chi^2 = \sum \frac{(O_i - E_i)^2}{E_i}, \quad df = (\text{num rows} - 1)(\text{num cols} - 1) \quad (79)$$

$$E_i = \frac{(\text{row total})(\text{column total})}{(\text{grand total})} = np_i \quad (80)$$

For 2×2 contingency tables, you can use the Fisher Exact Test:

fisher.test(D, alternative="greater")

(must specify alternative as greater)

9 Linear Regression

9.1 LINEAR CORRELATION

$H_0: \rho = 0$

cor.test(x, y)

where: **x** and **y** are ordered vectors.

$$r = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{(n-1)s_x s_y}, \quad t = \frac{r - 0}{\sqrt{\frac{1-r^2}{n-2}}}, \quad df = n - 2 \quad (81)$$

9.2 MODELS IN R

MODEL TYPE	EQUATION	R MODEL
linear 1 indep var	$y = b_0 + b_1 x_1$	$y \sim x_1$
... 0 intercept	$y = 0 + b_1 x_1$	$y \sim 0 + x_1$
linear 2 indep vars	$y = b_0 + b_1 x_1 + b_2 x_2$	$y \sim x_1 + x_2$
... interaction	$y = b_0 + b_1 x_1 + b_2 x_2 + b_{12} x_1 x_2$	$y \sim x_1 + x_2 + x_1 * x_2$
polynomial	$y = b_0 + b_1 x_1 + b_2 x_1^2$	$y \sim x_1 + 1(x_1^2)^2$

9.3 REGRESSION

Simple linear regression steps:

1. Make sure there is a significant linear correlation.
2. **results=lm(y~x)** Linear regression of y on x vectors
3. **results** View the results
4. **plot(x, y)**; **abline(results)** Plot regression line on data
5. **plot(x, results\$residuals)** Plot residuals

$$y = b_0 + b_1 x_1 \quad (82)$$

$$b_1 = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{\sum (x_i - \bar{x})^2} \quad (83)$$

$$b_0 = \bar{y} - b_1 \bar{x} \quad (84)$$

9.4 PREDICTION INTERVALS

To predict y when $x = 5$ and show the 95% prediction interval with regression model in results:

predict(results, newdata=data.frame(x=5), int="pred")

10 ANOVA

10.1 ONE WAY ANOVA

1. **results=aov(depVarColName~indepVarColName, data=tableName)** Run ANOVA with data in **tableName**, factor data in **indepVarColName** column, and response data in **depVarColName** column.
2. **summary(results)** Summarize results
3. **boxplot(depVarColName~indepVarColName, data=tableName)** Boxplot of levels for factor

$$F = \frac{MS(\text{treatment})}{MS(\text{error})}, \quad df_1 = k - 1, df_2 = N - k \quad (85)$$

To find required sample size and power see **power.anova.test(...)**

11 Loading and using external data and tables

11.1 LOADING EXCEL DATA

1. Export your table as a CSV file (comma separated file) from Excel.
2. Import your table into R using:
MyTable=read.csv(file.choose())

11.2 LOADING AN .RDATA FILE

You can either double click on the .RData file or use the menu:

- Windows: **File→Load Workspace...**
- Mac: **Workspace→Load Workspace File...**

11.3 USING TABLES OF DATA

1. To see all the available variables type: **ls()**
2. To see what's inside a variable, type its name.
3. If the variable **tableName** is a table, you can also type **names(tableName)** to see the column names or type **head(tableName)** to see the first few rows of data.
4. To access a column of data type **tableName\$columnName**

An example demonstrating how to get the women's height data and find the mean:

```
> ls() # See what variables are defined
[1] "women" "x"
> head(women) # Look at the first few entries
  height weight
1    58   115
2    59   117
3    60   120
> names(women) # Just get the column names
[1] "height" "weight"
> women$height # Display the height data
[1] 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72
> mean(women$height) # Find the mean of the heights
[1] 65
```